St. John Fisher University

Fisher Digital Publications

Pharmacy Faculty/Staff Publications

Wegmans School of Pharmacy

3-14-2013

Resolution of Clinical and Laboratory Abnormalities after Diagnosis of Ventilator-Associated Pneumonia in Trauma Patients

Joseph M. Swanson
Frederick Medical Center

Kathryn A. Connor
St. John Fisher University, kaconnor@sjf.edu

Louis J. Magnotti
University of Tennessee Health Services Center

Martin A. Croce
University of Tennessee Health Services Center

Jessica Johnson
Frederick Medical Center

Follow this and additional works at: https://fisherpub.sjf.edu/pharmacy_facpub
See next page for additional authors

Part of the [Pharmacy and Pharmaceutical Sciences Commons](#)

[How has open access to Fisher Digital Publications benefited you?](#)

Publication Information

Swanson, Joseph M.; Connor, Kathryn A.; Magnotti, Louis J.; Croce, Martin A.; Johnson, Jessica; Wood, G. Christopher; and Fabian, Timothy C. (2013). "Resolution of Clinical and Laboratory Abnormalities after Diagnosis of Ventilator-Associated Pneumonia in Trauma Patients." *Surgical Infections* 14.1, 49-55. Please note that the Publication Information provides general citation information and may not be appropriate for your discipline. To receive help in creating a citation based on your discipline, please visit <http://libguides.sjfc.edu/citations>.

This document is posted at https://fisherpub.sjf.edu/pharmacy_facpub/32 and is brought to you for free and open access by Fisher Digital Publications at . For more information, please contact fisherpub@sjf.edu.

Resolution of Clinical and Laboratory Abnormalities after Diagnosis of Ventilator-Associated Pneumonia in Trauma Patients

Abstract

Background: Guidelines advise that patients with ventilator-associated pneumonia (VAP) should respond clinically by Day 3 of antibiotics. White blood cell (WBC) count, maximum temperature (Tmax), and PaO₂:FIO₂ ratio are all said to respond significantly by Day 6. Resolution of abnormalities has not been evaluated in trauma patients.

Methods: Retrospective review of trauma patients with VAP. The WBC count, Tmax, and PaO₂:FIO₂ were evaluated for 16 days after diagnosis. Patients were grouped into uncomplicated VAP, complicated VAP (those with inadequate empirical therapy [IEAT], VAP relapse/superinfection, or acute respiratory distress syndrome), and concurrent infection +VAP (those also infected at another site).

Results: There were 126 patients (uncomplicated VAP = 29, complicated VAP = 69, and concurrent infection + VAP = 28). The mean Tmax in patients with uncomplicated VAP decreased significantly from diagnosis to Day 4 (Day 1: 39 – 0.5°C vs. Day 4: 38.6 – 0.7°C; $p = 0.028$) but never normalized. Their WBC counts and PaO₂:FIO₂ did not change significantly over the 16-day follow-up and never normalized. When comparing the three groups, the probability of resolving all three abnormalities was not different ($p = 0.5$).

Conclusions: Clinical and laboratory abnormalities in critically injured patients with VAP do not resolve as quickly as suggested in the guidelines. Future studies should evaluate new methods to determine the response to antibiotic therapy in critically injured patients with VAP.

Disciplines

Pharmacy and Pharmaceutical Sciences

Comments

This is a copy of an article published in Surgical Infections © 2013 [copyright Mary Ann Liebert, Inc.]; Surgical Infections is available online at: <http://online.liebertpub.com>.

Creative Commons License

This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Authors

Joseph M. Swanson, Kathryn A. Connor, Louis J. Magnotti, Martin A. Croce, Jessica Johnson, G. Christopher Wood, and Timothy C. Fabian

Resolution of Clinical and Laboratory Abnormalities after Diagnosis of Ventilator-Associated Pneumonia in Trauma Patients

Joseph M. Swanson,¹ Kathryn A. Connor,² Louis J. Magnotti,³ Martin A. Croce,³
Jessica Johnson,¹ G. Christopher Wood,¹ and Timothy C. Fabian³

Abstract

Background: Guidelines advise that patients with ventilator-associated pneumonia (VAP) should respond clinically by Day 3 of antibiotics. White blood cell (WBC) count, maximum temperature ($T_{\max}$), and $\text{PaO}_2\text{:F}_i\text{O}_2$ ratio are all said to respond significantly by Day 6. Resolution of abnormalities has not been evaluated in trauma patients.

Methods: Retrospective review of trauma patients with VAP. The WBC count, $T_{\max}$, and $\text{PaO}_2\text{:F}_i\text{O}_2$ were evaluated for 16 days after diagnosis. Patients were grouped into uncomplicated VAP, complicated VAP (those with inadequate empirical therapy [IEAT], VAP relapse/superinfection, or acute respiratory distress syndrome), and concurrent infection + VAP (those also infected at another site).

Results: There were 126 patients (uncomplicated VAP=29, complicated VAP=69, and concurrent infection + VAP=28). The mean $T_{\max}$ in patients with uncomplicated VAP decreased significantly from diagnosis to Day 4 (Day 1: $39 \pm 0.5^\circ\text{C}$ vs. Day 4: $38.6 \pm 0.7^\circ\text{C}$; $p=0.028$) but never normalized. Their WBC counts and $\text{PaO}_2\text{:F}_i\text{O}_2$ did not change significantly over the 16-day follow-up and never normalized. When comparing the three groups, the probability of resolving all three abnormalities was not different ($p=0.5$).

Conclusions: Clinical and laboratory abnormalities in critically injured patients with VAP do not resolve as quickly as suggested in the guidelines. Future studies should evaluate new methods to determine the response to antibiotic therapy in critically injured patients with VAP.

VENTILATOR-ASSOCIATED PNEUMONIA (VAP) is the most common serious infection in critically ill, mechanically ventilated patients and carries a high mortality rate (33%–50%) [1]. It is essential to treat VAP appropriately with antibiotics and evaluate a patient's response to therapy by monitoring for improvement or normalization of clinical signs, including temperature, white blood cell (WBC) count, and $\text{PaO}_2\text{:F}_i\text{O}_2$ ratio. Such monitoring is crucial because non-response may indicate suboptimal treatment or a drug-resistant pathogen. Current guidelines advise that appropriately treated patients should have a good clinical response by Day 3 and a substantive improvement by Day 6 [1]. These recommendations are based mainly on expert opinion and two studies [2,3] in which clinical improvement occurred by Days 3–5, as judged primarily by WBC count, maximum temperature ($T_{\max}$), and $\text{PaO}_2\text{:F}_i\text{O}_2$. Using these guidelines, some practitioners initiate empirical antibiotics for clinical

suspicion of VAP and then use a good clinical response to confirm the diagnosis.

However, trauma patients are a unique population that may respond to VAP treatment differently than do general medical/surgical intensive care unit (ICU) patients. Traumatic injury can induce the systemic inflammatory response syndrome (SIRS), contributing to derangements of clinical measures used to determine the response to VAP therapy. As many as 91% of trauma patients exhibit signs of SIRS during the first week in the ICU [4]. In addition, a composite score of clinical measures (Clinical Pulmonary Infection Score [CPIS]) used for VAP diagnosis in some medical ICUs does not perform well in critically injured patients, making it a poor choice for evaluating the response to therapy [5].

We hypothesized that clinical and laboratory abnormalities in patients with no confounders would decline significantly after VAP, whereas those in patients with confounders would

¹Department of Pharmacy, Frederick Medical Center, Frederick, Maryland.

²St. John Fisher College, University of Rochester Medical Center, Rochester, New York.

³Department of Surgery, University of Tennessee Health Sciences Center, Memphis, Tennessee.

Presented at the 39th Critical Care Congress Society of Critical Care Medicine, Miami, Florida, January 28–31, 2010.

stay elevated throughout follow-up. Additionally, because there are insufficient data to characterize the clinical and laboratory response to VAP in critically injured patients, we evaluated the resolution of clinical and laboratory abnormalities retrospectively in these patients after diagnosis and appropriate treatment of VAP.

Patients and Methods

Following approval by the University of Tennessee Institutional Review Board (IRB number 09-00277-XM), this retrospective review was conducted at the Presley Memorial Trauma Center, Memphis, TN. Patients admitted from December 1, 2006, through July 31, 2008, were eligible if they were ≥ 18 years of age and had VAP diagnosed. Patients who died during treatment were excluded, as their elevated values prior to death could mask overall resolution in the group. Patients receiving steroids also were excluded in order to avoid misinterpretation of $T_{\max}$ and WBC count.

Diagnosis and treatment of VAP

Ventilator-associated pneumonia was suspected if patients had a new or changing infiltrate on chest radiograph and experienced two of the following clinical signs: fever or hypothermia ($>38^{\circ}\text{C}$ or $<36^{\circ}\text{C}$), leukocytosis or leukopenia ($>10 \times 10^3$ cells/ mm^3 or $<4 \times 10^3$ cells/ mm^3), or macroscopically purulent sputum. The VAP diagnosis was confirmed using bronchoscopic bronchoalveolar lavage (BAL) with quantitative culture demonstrating bacterial growth $\geq 100,000$ colony-forming units (CFU)/mL. Empirical antibiotic therapy was initiated in all patients immediately after the BAL. All patients were treated for suspected VAP using a unit-specific protocol [5,6–8]. Thus, patients admitted to the trauma intensive care unit (TICU) for ≤ 7 days received ampicillin/sulbactam 3 g intravenously (IV) every 6 h, or moxifloxacin 400 mg IV every 24 h in those with penicillin allergy. Patients mechanically ventilated for ≥ 7 days received vancomycin 20 mg/kg IV q 12 h plus cefepime 2 g IV q 8 h, or ciprofloxacin 400 mg IV q 8 h for those with penicillin allergy. On identification of the causative pathogen, antibiotic therapy was narrowed appropriately. Based on previous data, the treatment protocol dictates that patients with early VAP (<7 days) be treated for seven days [7]. For late VAP, the protocol recommends a follow-up BAL to determine microbiologic response to antibiotic therapy, if possible. A follow-up BAL demonstrating growth of the original pathogen of $<10,000$ CFU/mL is deemed a microbiologic response. If growth is 10,000–99,999 CFU/mL, antibiotics are continued for the maximum duration to ensure appropriate treatment. If culture of an organism yields $\geq 100,000$ CFU/mL, the BAL finding is deemed a relapse or superinfection (depending on the original pathogen). The protocol recommends a maximum duration of therapy of 14 days for late pathogens such as methicillin-resistant *Staphylococcus aureus* (MRSA), *Pseudomonas aeruginosa*, or *Acinetobacter baumannii* [9]. If patients' signs and symptoms are not resolving while the patient is being treated, the protocol recommends continuing to search for new or recurrent infections as indicated (subsequent BALs, blood cultures, urine cultures, etc.). If the signs of VAP are not resolved at the end of antibiotic therapy, antibiotics are stopped, and the search for occult infections continues. Extreme cases such as hemodynamic instability could warrant

continuation of antibiotics, but this is not common. This approach is supported by repeat microbiologic culture data from our institution [7,9].

Study groups

All patients experienced VAP, were treated with appropriate definitive antibiotics, and survived the VAP episode. Patients were categorized into one of three groups based on complicating factors that could affect the resolution of clinical evidence of disease. Patients with VAP who were treated appropriately, did not acquire any other infections, and did not have any of the complicating factors listed below were placed in the uncomplicated VAP group. Patients with complications that could relate to VAP, such as inadequate empiric antibiotic therapy (IEAT), VAP relapse, VAP superinfection, or acute respiratory distress syndrome (ARDS), were placed in the complicated VAP group. Patients with concurrent infection(s) at another site were placed in the concurrent infection + VAP group.

Definitions

The adequacy of empiric therapy was determined for each VAP episode. IEAT was defined as VAP caused by at least one bacterial species reported to have either intermediate or full in vitro resistance to all empirical antibiotics. Empiric antibiotic therapy was considered adequate if at least one antibiotic in the regimen demonstrated in vitro activity against all identified pathogens. The treating physician diagnosed ARDS using standard criteria that included acute onset, bilateral infiltrates on chest radiograph, exclusion of cardiogenic pulmonary edema, and a $\text{PaO}_2:\text{FiO}_2 < 200$. Additional infections were defined as follows: VAP relapse, a subsequent BAL demonstrating growth of the original pathogen $\geq 100,000$ CFU/mL; VAP superinfection, a subsequent BAL demonstrating growth of a different pathogen $\geq 100,000$ CFU/mL, bacteremia/fungemia, growth from at least two bottles for *S. epidermidis* or from one bottle for all other bacteria or fungi; urinary tract infection (UTI), urine culture demonstrating bacterial or fungal growth $\geq 100,000$ CFU/mL; surgical site infection, or any wound deemed infected by the treating surgeon and requiring antibiotic therapy.

Data collection

Demographic data, Injury Severity Score (ISS), duration of mechanical ventilation, and length of stay in the TICU and hospital were recorded. Patients were followed from the day the BAL was conducted (Day 1) through Day 16. Patients who were discharged from the ICU during the follow-up period were included in the final analysis. Patients discharged from the hospital are clinically stable. Therefore, unresolved clinical abnormalities were considered resolved on the day of discharge and were categorized as such through Day 16. The following clinical and laboratory data were recorded daily: $T_{\max}$, WBC count, and $\text{PaO}_2:\text{FiO}_2$. Resolution was defined as follows: $T_{\max} \geq 36^{\circ}\text{C}$ or $\leq 38^{\circ}\text{C}$, WBC count $\geq 4 \times 10^3$ cells/ mm^3 or $\leq 10 \times 10^3$ cells/ mm^3 , and a $\text{PaO}_2:\text{FiO}_2$ ratio ≥ 250 [2,10]. The diagnosis of VAP does not require all disease features be abnormal; thus, resolution for individual measures was evaluated only in those patients with abnormal variables on the day of diagnosis.

Statistical analysis

Continuous variables were compared using the Student *t*-test for normally distributed variables or the Wilcoxon rank-sum test for non-normally distributed variables. Proportions were compared using the χ^2 test. Repeated-measures analysis of variance was used to compare changes in clinical features over the 16-day follow-up period for the three groups. Multiple pairwise comparisons were made using the Holm-Sidak method. The Kaplan-Meier method was used to depict the probability of resolution of clinical and laboratory abnormalities. The log-rank test was used to compare two Kaplan-Meier curves. All statistics were calculated using SigmaPlot 11.1 software (Systat Software, Inc., San Jose, CA). Data are presented as mean $\pm$ standard deviation (SD), mean $\pm$ 95% confidence interval (95% CI), median (interquartile range), or proportions, as indicated.

Results

A total of 142 patients with VAP were identified. Fifteen patients were excluded because of death during antibiotic therapy for VAP, and one was excluded because insufficient information was available. Therefore, 126 patients were analyzed. Twenty-nine patients (23%) had no complications during their VAP episode (uncomplicated VAP), whereas 69 patients (54.8%) experienced IEAT, a VAP relapse, superinfection, or ARDS during their VAP episode (complicated VAP). Twenty-eight patients (22.2%) had a concurrent UTI, bacteremia, or surgical site infection during their VAP episode (concurrent infection+VAP). Patient demographics were similar in all three groups (Table 1). It was not surprising that patients with complicated VAP had a significantly lower rate of early VAP ($p=0.006$) and a statistical trend toward a higher ISS ($p=0.057$).

The VAP pathogens are listed in Table 2. Of note, *Haemophilus influenzae* was the most common. Methicillin-resistant

S. aureus accounted for 7.4% of organisms overall, 7.4% in the uncomplicated VAP group, 6.7% in the complicated VAP group, and 9.5% in the concurrent infection + VAP group.

Overall, 90 patients (71%) underwent one surgical procedure during the 16-day follow-up. This first operation was performed on average 5.4 ± 4.5 days (range 1–15 days) after the diagnosis of VAP. Thirty-nine patients (31%) underwent a second operation 7.4 ± 4.1 days after VAP diagnosis, and smaller numbers of patients underwent three, four, and five surgical procedures (14, 7, and 3 patients, respectively). Operation 1 occurred in 20 patients (69%) with uncomplicated VAP, 51 (74%) with complicated VAP, and 19 (68%) with concurrent infections+VAP. Operation 2 occurred in 10 patients (35%) with uncomplicated VAP, 23 (33%) of those with complicated VAP, and six patients (21%) with concurrent infections + VAP.

Changes in clinical and laboratory measures for all three groups are depicted in Figure 1. In patients with uncomplicated VAP and an elevated temperature at baseline, the earliest statistically significant decrease in $T_{\max}$ occurred on Day 4 (Day 1: $39 \pm 0.5^\circ\text{C}$ vs. Day 4: $38.6 \pm 0.7^\circ\text{C}$, $p=0.028$). However, the group's mean $T_{\max}$ never dropped below 38°C (Day 16 mean $T_{\max}$ $38.1 \pm 0.6^\circ\text{C}$). Patients with complicated VAP with an elevated baseline temperature demonstrated a statistically significant decrease in $T_{\max}$ at Day 2 (Day 1: $39.1 \pm 0.5^\circ\text{C}$ vs. Day 2: $38.7 \pm 0.8^\circ\text{C}$; $p=0.047$). Interestingly, the mean $T_{\max}$ normalized on Day 16 in this group (mean $38 \pm 0.8^\circ\text{C}$). In the concurrent infection+VAP group, $T_{\max}$ decreased significantly on Day 7 (Day 1: $39.2 \pm 0.5^\circ\text{C}$ vs. Day 7: $38.5 \pm 0.7^\circ\text{C}$; $p=0.005$). The mean $T_{\max}$ normalized on Day 13 (mean $38 \pm 0.8^\circ\text{C}$), but then increased again and normalized on Day 16 (mean $37.9 \pm 0.8^\circ\text{C}$).

In patients with an elevated baseline WBC count, there was no statistically significant change over the 16-day follow-up (uncomplicated VAP $p=0.26$; Complicated VAP $p=0.097$; concurrent infection+VAP $p=0.09$), and the mean WBC

TABLE 1. DEMOGRAPHICS OF STUDY PATIENTS BY GROUP

	Uncomplicated VAP (n=29)	Complicated VAP (n=69)	Concurrent infection with VAP (n=28)	p value ^a
Mean age (years $\pm$ SD)	44.2 $\pm$ 15.8	42.8 $\pm$ 17.2	44.6 $\pm$ 19.3	NS
Male/female	22/7	56/13	21/7	NS
Blunt injury, % (n)	86 (25)	84 (58)	75 (21)	NS
Type of injury, % (n)				
TBI	42 (12)	55 (38)	54 (15)	NS
SCI	7 (2)	13 (9)	7 (2)	NS
Fractures	79 (23)	68 (47)	54 (15)	NS
Abdominal	35 (10)	43 (30)	43 (12)	NS
Chest	45 (13)	45 (31)	54 (15)	NS
Median Injury Severity Score (IQR)	26 (20–34)	34 (25–42)	31 (22–41)	NS
Early VAP, % (n)	79 (23)	46 (32)	68 (19)	0.006
ARDS, % (n)	0 (0)	42 (29)	0 (0)	
Subsequent BAL, % (n)	62 (18)	84 (58)	54 (15)	
Microbiologic response on BAL, % (n)	100 (18)	52 (30)	100 (15)	
Median days of mechanical ventilation (IQR)	17 (13–33)	24 (16–47)	18 (11–29)	NS
Median ICU length of stay (IQR)	26 (18–42)	28 (20–49)	21 (16–36)	NS
Median hospital length of stay (IQR)	35 (26–52)	36 (26–61)	33 (29–41)	NS

^aComparison between VAP only and VAP+confounders.

ARDS=acute respiratory distress syndrome; BAL=bronchoalveolar lavage; ICU=intensive care unit; IQR=interquartile range; SCI=spinal cord injury; SD=standard deviation; TBI=traumatic brain injury; VAP=ventilator-associated pneumonia.

TABLE 2. PATHOGENS ISOLATED FROM PATIENTS WITH VENTILATOR-ASSOCIATED PNEUMONIA

Pathogen	Uncomplicated VAP (n = 29)	Complicated VAP (n = 68)	Concurrent infection with VAP (n = 28)	Total (n = 126)
Gram-positive organisms				
Methicillin-sensitive	8	12	6	26
Staphylococcus aureus				
Methicillin-resistant S. aureus	4	8	4	16
Viridans Streptococcus	6	10	2	18
Streptococcus pneumoniae	4	5	5	14
Other streptococci	1	3	2	6
Methicillin-sensitive S. epidermidis	0	1	0	1
Methicillin-resistant S. epidermidis	0	1	0	1
Diphtheroids	1	3	1	5
Gram-negative organisms				
Citrobacter koseri	1	0	0	1
Haemophilus influenzae	9	21	9	39
Haemophilus parainfluenzae	3	4	0	7
Acinetobacter baumannii	0	12	6	18
Pseudomonas aeruginosa	3	11	2	16
Proteus mirabilis	1	2	1	4
Escherichia coli	3	2	0	5
Enterobacter aerogenes	2	4	0	6
Enterobacter cloacae	0	3	0	3
Non-pathogenic Neisseria species	4	2	2	8
Achromabacter species	0	2	0	2
Aeromonas hydrophilia	0	1	0	1
Stenotrophomonas maltophilia	0	8	0	8
Serratia marcescens	0	2	0	2
Klebsiella pneumoniae	1	3	1	5
Branhamella catarrhalis	3	0	1	4
Total organisms	54	120	42	216

count never normalized in any of the three groups. The $\text{PaO}_2\text{:F}_i\text{O}_2$ in the uncomplicated VAP and concurrent infection + VAP groups did not change significantly ($p=0.058$ and 0.197 , respectively), whereas the complicated VAP group demonstrated a significant decrease at Day 3 (Day 1: 170 ± 67 vs. Day 3: 246 ± 158 ; $p=0.033$). The $\text{PaO}_2\text{:F}_i\text{O}_2$ fluctuated by worsening on Days 4–6, but improving significantly on Days 8, 9, 13, and 16.

The proportions of patients whose clinical and laboratory abnormalities resolved on Days 3, 6, 9, 12, and 16 are presented in Table 3. The majority of patients in all three groups did not achieve resolution by Day 6. Even by Day 16, resolution was less common than had been expected. The probability of resolving all three abnormalities is displayed in Figure 2. Kaplan-Meier log-rank analysis demonstrated no significant difference between patients with uncomplicated VAP, complicated VAP, and concurrent infection + VAP. In addition, similar results were seen when patients were categorized according to the timing of VAP (early or late) or whether the VAP was caused by non-fermenting gram-negative bacilli (data not shown).

Discussion

This retrospective study raises serious concerns about the value of the traditional measures used to assess VAP treatment in trauma patients. According to the current American Thoracic Society/Infectious Diseases Society of America (ATS/IDSA) guidelines [1], the uncomplicated VAP group

should have demonstrated a clear response to therapy by Day 3 and a substantive response by Day 6. However, in this group, the T_{max} was the only measure to decline significantly during the 16-day follow-up. Although the decline was statistically significant, it was likely not clinically significant, being approximately 0.4°C by Day 4. Indeed, the group mean never normalized. The WBC count and $\text{PaO}_2\text{:F}_i\text{O}_2$ never declined significantly and never normalized. These findings seem valid because the uncomplicated VAP group had no concomitant infections or complicating factors related to VAP that may have kept the clinical and laboratory abnormalities from resolving. Also, all patients in this group were confirmed to have had a positive microbiologic response to treatment according to a follow-up BAL. Furthermore, the clinical response did not differentiate patients with uncomplicated VAP, complicated VAP, or concurrent infection + VAP. Traditional medical belief dictates that there probably would have been a clear difference in the resolution of the clinical and laboratory abnormalities in these groups. Unfortunately, in these critically injured patients, resolution of all disease measures was poor in each of the three groups (less than 35% by Day 16) despite successful treatment of their VAP. Additionally, the probability of resolution of all three measures did not differ between groups. Overall, these results suggest that the current guidelines do not apply well to trauma patients.

There are only a few studies that have evaluated this issue. The study most similar in design to the current one was performed by Dennesen et al. [2], but their results were different from ours. They noted T_{max} and WBC count declined and

FIG. 1. Mean values for clinical and laboratory measures after diagnosis of ventilator-associated pneumonia. Day 1 is the day bronchoalveolar lavage was conducted to verify the pneumonia.

the $\text{PaO}_2:\text{FiO}_2$ increased after VAP. However, it is important to note that they enrolled medical, surgical, trauma, and neurologic patients, whereas the current study included only trauma patients. In addition, in the current study, the WBC count and T_{max} were higher, and the $\text{PaO}_2:\text{FiO}_2$ was lower, on the day of VAP diagnosis than in the study by Dennesen et al. (WBC $15.1 \pm 3.8 \times 10^3$ cells/ mm^3 vs. $14.2 \pm 5.4 \times 10^3$ cells/ mm^3 , T_{max} $39 \pm 0.6^\circ\text{C}$ vs. $38.8 \pm 0.5^\circ\text{C}$, $\text{PaO}_2:\text{FiO}_2$ 176 ± 55 vs. 197 ± 79). Perhaps the patients in the current study were more seriously ill at the onset of infection. Moreover, Dennesen

et al. used a resolution cutoff for the $\text{PaO}_2:\text{FiO}_2$ of 188, which is lower than that used in the current study and by others [10,11]. Another difference is that the diagnostic criteria for VAP in the current study were more rigorous. Dennesen et al. did not define their diagnostic threshold, whereas the current study used $\geq 100,000$ CFU/mL, which is recommended in the ATS/IDSA guidelines but is not accepted universally. Finally, the paper by Dennesen et al. is somewhat confusing because they report a mean duration until resolution of six days for all three signs, whereas their graph indicates that the mean WBC count never normalized.

Other studies were similar to the current study in showing that clinical and laboratory abnormalities do not resolve quickly or cleanly during VAP. A large randomized trial [12] noted only a small decline in mean temperature after treatment of VAP (Day 1 $\sim 38.6^\circ\text{C}$, Day 2 $\sim 38.1^\circ\text{C}$). Although the temperature responded to antibiotic therapy within three days, the mean $\text{PaO}_2:\text{FiO}_2$ did not increase above 250 until after Day 7, and the mean WBC count remained above 10×10^3 cells/mL through Day 28.

A related issue is the possibility that IEAT or specific pathogens can affect resolution of disease. Vidaur et al. reported the clinical resolution of 60 patients with adequate empiric therapy and 30 with inadequate therapy [10]. Inadequate EAT delayed the median time to resolution of fever, hypoxemia, and WBC count by 5–7 days, 6–8 days, and 3–5 days, respectively. Infection with MRSA increased the time to resolution of disease regardless of IEAT status. These data highlight additional factors that must be considered when evaluating the treatment response. In the current study, patients experiencing IEAT were separated from those receiving appropriate therapy. Patients with MRSA VAP were not grouped separately because the incidence of MRSA was similar in the three groups (uncomplicated VAP 7.4%, complicated VAP 6.7%, and concurrent infection + VAP 9.5%). Thus, these factors are unlikely to explain the delayed response seen in the current study.

The lack of resolution in the current study likely is multifactorial. Surgery has been associated with elevations in temperature and WBC count even if the patient does not have an infection [13]. Thus, the multiple surgical procedures performed in the study patients may have masked a clinical response to antibiotics by altering temperature and WBC count. Additionally, more than one-half of the patients had early VAP (< 7 days), suggesting the presence of SIRS as a result of their original trauma [4].

New methods must be identified for evaluating the response of trauma patients to VAP therapy. Repeat BAL cultures can be used [7,14] and were the standard of practice in the current study. Unlike clinical measures, repeat cultures are not affected by non-infectious variables and thus are a more objective marker. Mueller et al. noted shorter durations of antibiotic therapy when repeat BAL cultures were used [7]. Microbiologic data could reduce clinicians' reliance on other laboratory data, especially in trauma patients. However, repeat cultures are invasive and can be expensive, depending on the techniques used. Alternatively, biomarkers such as procalcitonin may be able to signal response to therapy [15], and in one study this protein was not affected by surgery [16]. A recent meta-analysis of seven studies showed that procalcitonin monitoring decreased the duration of antibiotics in patients with respiratory tract infections [15].

TABLE 3. RESOLUTION OF CLINICAL ABNORMALITIES

Day of resolution	Uncomplicated VAP	Complicated VAP	Concurrent infection + VAP
Maximum temperature, % (n/total n)			
Day 3	10.3 (3/29)	18 (11/61)	4.2 (1/24)
Day 6	27.6 (5/29)	24.6 (15/61)	16.7 (4/24)
Day 9	37.9 (11/29)	32.8 (20/61)	37.5 (9/24)
Day 12	55.2 (16/29)	42.6 (26/61)	37.5 (9/24)
Day 16	51.7 (15/29)	53.4 (35/61)	54.2 (13/24)
White blood cell count, % (n/total n)			
Day 3	21.1 (4/19)	18.9 (7/37)	10.5 (2/19)
Day 6	21.1 (4/19)	10.8 (4/37)	31.6 (6/19)
Day 9	15.8 (3/19)	10.8 (4/37)	31.6 (6/19)
Day 12	26.3 (5/19)	16.2 (6/37)	15.8 (3/19)
Day 16	15.8 (3/19)	35.1 (13/37)	36.8 (7/19)
PaO ₂ :F _i O ₂ , % (n/total n)			
Day 3	29.2 (7/24)	33.9 (19/56)	40.9 (9/22)
Day 6	45.8 (11/24)	30.4 (17/56)	40.9 (9/22)
Day 9	33.3 (8/24)	41.1(23/56)	59.1 (13/22)
Day 12	29.2 (7/24)	51.8 (29/56)	40.9 (9/22)
Day 16	33.3 (8/24)	62.5 (35/56)	72.7 (16/22)
All abnormalities, % (n/total n)			
Day 3	3.4 (1/29)	4.3 (3/69)	3.5 (1/28)
Day 6	10.3 (3/29)	4.3 (3/69)	0 (0/28)
Day 9	10.3 (3/29)	4.3 (3/69)	3.6 (1/28)
Day 12	10.3 (3/29)	7.2 (5/69)	7.1 (2/28)
Day 16	20.1 (6/29)	21.7 (15/69)	32.1 (9/28)

The current study has several limitations. The data were collected retrospectively, which is inherently less specific than a prospective study. However, the variables were recorded accurately and would not have been different if collected prospectively. Second, a diagnostic threshold for VAP

of $\geq 100,000$ CFU/mL is higher than the $\geq 10,000$ CFU/mL used by many institutions. The larger bacterial load may result in a longer time to resolution of clinical and laboratory abnormalities. Finally, the study enrolled only 29 patients in the uncomplicated VAP group, which is the ideal study group for this problem. Although this number is relatively small, it is not substantially different from the enrollment in other published studies evaluating resolution of disease [2,3]. Indeed, the total number of patients across the three groups is good in relation to this body of literature. The strengths of the study include a relatively homogeneous population, the use of the highest-quality VAP diagnostic criteria, and a high percentage of patients with follow-up cultures that confirmed a positive response to treatment. The study enrolled only critically injured patients, a population that is inherently different from other critically ill patients. Use of bronchoscopic BAL with quantitative cultures ensured that the patients actually had VAP and thus eliminates bias from including patients without pneumonia.

Conclusion

Signs, symptoms, and laboratory abnormalities in critically injured trauma patients with VAP do not resolve as quickly as suggested by the guidelines. The probability of resolving all three clinical measures does not differ in patients with uncomplicated VAP and those with complications or concurrent infections. Future studies should evaluate new methods to determine the response to antibiotic therapy in critically injured patients with VAP.

Author Disclosure Statement

No competing financial interests exist for any of the authors.

FIG. 2. Kaplan-Meier graph depicting the probability that all three abnormalities (maximum temperature, white blood cell count, and PaO₂:F_iO₂) will not resolve over the 16-day follow-up period. There was no difference in the probability of non-resolution ($p=0.5$).

References

- Guidelines for the management of adults with hospital-acquired, ventilator-associated, and healthcare-associated pneumonia. *Am J Respir Crit Care Med* 2005;171:388–416.
- Dennesen PJ, van der Ven AJ, Kessels AG, et al. Resolution of infectious parameters after antimicrobial therapy in patients with ventilator-associated pneumonia. *Am J Respir Crit Care Med* 2001;163:1371–1375.
- Luna CM, Blanzaco D, Niederman MS, et al. Resolution of ventilator-associated pneumonia: Prospective evaluation of the clinical pulmonary infection score as an early clinical predictor of outcome. *Crit Care Med* 2003;31:676–682.
- Hoover L, Bochicchio GV, Napolitano LM, et al. Systemic inflammatory response syndrome and nosocomial infection in trauma. *J Trauma* 2006;61:310–316.
- Croce MA, Swanson JM, Magnotti LJ, et al. The futility of the clinical pulmonary infection score in trauma patients. *J Trauma* 2006;60:523–527.
- Croce MA, Fabian TC, Mueller EW, et al. The appropriate diagnostic threshold for ventilator-associated pneumonia using quantitative cultures. *J Trauma* 2004;56:931–934.
- Mueller EW, Croce MA, Boucher BA, et al. Repeat bronchoalveolar lavage to guide antibiotic duration for ventilator-associated pneumonia. *J Trauma* 2007;63:1329–1337.
- Swanson JM, Wood GC, Croce MA, et al. Utility of preliminary bronchoalveolar lavage results in suspected ventilator-associated pneumonia. *J Trauma* 2008;65:1271–1277.
- Magnotti LJ, Croce MA, Zarzaur BL, et al. Causative pathogen dictates optimal duration of antimicrobial therapy for ventilator-associated pneumonia in trauma patients. *J Am Coll Surg* 2011;212:476–484.
- Vidaur L, Planas K, Sierra R, et al. Ventilator-associated pneumonia: Impact of organisms on clinical resolution and medical resources utilization. *Chest* 2008;133:625–632.
- Myrianthefs P, Ioannides K, Gavala A, et al. Patterns of resolution of infectious parameters in multidrug-resistant *Acinetobacter baumannii* ventilator-associated pneumonia. *J Hosp Infect* 2010;75:79–80.
- Chastre J, Wolff M, Fagon JY, et al. Comparison of 8 vs. 15 days of antibiotic therapy for ventilator-associated pneumonia in adults: A randomized trial. *JAMA* 2003;290:2588–2598.
- Kumar S, Mehta Y, Vats M, et al. An observational study to know the association of leukocytosis and fever with infection in post cardiac surgery patients. *Indian Heart J* 2007;59:316–322.
- Wunderink RG, Mendelson MH, Somero MS, et al. Early microbiological response to linezolid vs. vancomycin in ventilator-associated pneumonia due to methicillin-resistant *Staphylococcus aureus*. *Chest* 2008;134:1200–1207.
- Li H, Luo YF, Blackwell TS, Xie CM. Meta-analysis and systematic review of procalcitonin-guided therapy in respiratory tract infections. *Antimicrob Agents Chemother* 2011;55:5900–5906.
- Hunziker S, Hugle T, Schuchardt K, et al. The value of serum procalcitonin level for differentiation of infectious from noninfectious causes of fever after orthopaedic surgery. *J Bone Joint Surg Am* 2010;92:138–148.

Address correspondence to:

Dr. Joseph M. Swanson

University of Tennessee Health Sciences Center

881 Madison Ave., Room 203

Memphis, TN 38163

E-mail: jswanson@uthsc.edu